

# BIOLOGICAL CONTROL OF POSTHARVEST DISEASES OF FRUITS

---

Wojciech J. Janisiewicz<sup>1</sup> and Lise Korsten<sup>2</sup>

<sup>1</sup>*Appalachian Fruit Research Station, USDA Agricultural Research Service, Kearneysville, West Virginia 25430; e-mail: wjanisie@afrs.ars.usda.gov*

<sup>2</sup>*Department of Microbiology and Plant Pathology, University of Pretoria, Pretoria 0002, Republic of South Africa; e-mail: lkorsten@fabi.up.ac.za*

**Key Words** fruit decay, biocontrol, microbial interaction, microbial antagonists

■ **Abstract** Losses from postharvest fruit diseases range from 1 to 20 percent in the United States, depending on the commodity. The application of fungicides to fruits after harvest to reduce decay has been increasingly curtailed by the development of pathogen resistance to many key fungicides, the lack of replacement fungicides, negative public perception regarding the safety of pesticides and consequent restrictions on fungicide use. Biological control of postharvest diseases (BCPD) has emerged as an effective alternative. Because wound-invading necrotrophic pathogens are vulnerable to biocontrol, antagonists can be applied directly to the targeted area (fruit wounds), and a single application using existing delivery systems (drenches, line sprayers, on-line dips) can significantly reduce fruit decays. The pioneering biocontrol products BioSave and Aspire were registered by EPA in 1995 for control of postharvest rots of pome and citrus fruit, respectively, and are commercially available. The limitations of these biocontrol products can be addressed by enhancing biocontrol through manipulation of the environment, using mixtures of beneficial organisms, physiological and genetic enhancement of the biocontrol mechanisms, manipulation of formulations, and integration of biocontrol with other alternative methods that alone do not provide adequate protection but in combination with biocontrol provide additive or synergistic effects.

## INTRODUCTION

The battle against postharvest decays of fruits and vegetables has been fought for decades but has not yet been won. Even the average consumer, who shops for quality fresh fruits and vegetables and must often discard spoiled produce, recognizes the persistent problem of postharvest decay. Although the development of modern fungicides and improved storage technologies in the 1960s and 1970s have greatly extended the shelf life of fruit after harvest, postharvest losses vary from an estimated 5 percent to more than 20 percent in the United States, depending on the commodity (24), and can be as high as 50 percent in developing countries (52). Postharvest losses have been reduced mainly through postharvest fungicides

(52, 53) and, to a lesser degree, through postharvest management practices to reduce inoculum or effective management of the cold chain system (keeping produce at low temperatures, which greatly reduce pathogen growth, from harvest to retail). However, postharvest use of fungicides has been increasingly curtailed by the development of pathogen resistance to many key fungicides (74, 137, 148, 180), lack of replacement fungicides, and public perception that pesticides are harmful to human health and the environment. This negative perception has promoted governmental policies restricting use of fungicides (73, 143). Thus, alternative methods to control postharvest diseases are urgently needed (35, 60, 157, 158, 165, 177, 183). Over the past 15 years, biological control has emerged as an effective strategy to combat major postharvest decays of fruits (82, 86, 112, 186). Compared to the long-standing interest in biological control of soilborne pathogens (182), research into biological control of postharvest decays (BCPD) is in its infancy. Nevertheless, progress has been substantial; the first commercial products have been registered in the United States by the U.S. Environmental Protection Agency (EPA) and are sold under the names BioSave100 and 110 and Aspire. In other countries such as South Africa, biocontrol products for the control of fruit diseases are registered [National Department of Agriculture Fertilizer, Farm Feeds, Agricultural and Stock Remedies (Act 36 of 1947)], and sold as Avogreen and YieldPlus.

Over two dozen programs worldwide are currently under way to develop BCPD of fruits, encompassing new approaches and methods to fit the postharvest system. General protocols for efficient discovery, scale-up, and pilot testing, necessary for rapid progress in developing biocontrol (37) have been developed and are driving this burgeoning area of research. In Europe, a multinational project is operating under the auspices of the European Commission (<http://www.biopostharvest.org/>). Commercial products are in the advanced stages of development and should reach European markets within the next year or two. Within governmental and industrial circles, there is firm commitment to accelerate development of new alternative strategies for the control of postharvest diseases.

This review describes the uniqueness and key strategies underlying the development of biological control of postharvest diseases of fruits, possible approaches to increase the spectrum of activity, commercial successes, and prospects for the future.

## UNIQUENESS OF THE POSTHARVEST SYSTEM

Progress in BCPD, especially in the postharvest application of antagonists, may be attributed to the uniqueness and relative simplicity of the postharvest system. Wounds made during harvesting and fruit handling can be protected from wound-invading pathogens with a single postharvest application of the antagonist directly to wounds, using existing delivery systems (drenches, on-line sprayers, on-line dips). Once harvested, fruits are placed in cold storage for various periods of time

ranging from a few days to months, depending on the commodity. Environmental conditions, such as temperature and relative humidity, can be managed to favor antagonist survival. Furthermore, biotic interference is minimal so antagonists encounter minimal competition from indigenous microorganisms. Consequently, biocontrol of postharvest diseases tends to be more consistent than biocontrol under field conditions, and the occasional variation in performance usually can be traced to nonstandard procedures or conditions. The levels of decay acceptable in postharvest systems are generally below 5 percent, a standard that is often achieved with a single application of the biocontrol agents. Given the high retail value of fresh fruits, the application of high concentrations of the antagonists to fruit surfaces is economically viable, whereas under field conditions this usage might not be cost effective.

The short period between harvesting and placing fruit in storage, from less than a day to a few days, requires rapid antagonist action. Once fruit is placed in cold storage, metabolic rates of the host and associated microflora will decline depending on the temperature regime selected. The search for antagonists to control postharvest wound invading pathogens should be narrowed to rapid colonizers of the wound site that can still be metabolically active at low storage temperatures.

The ecology of the targeted pathogen must be understood in developing a biological control strategy. Pathogens that tolerate environmental stress often have few competitors, since few species can exist under such conditions. For example, the opportunistic pathogen *Botrytis cinerea* may be a poor competitor (14) in comparison to *Penicillium* spp., which often produce secondary metabolites that inhibit competitors. Stress-tolerant and competitive species would therefore require biocontrol strategies different from those of ruderal species (14), which depend upon physical adaptations to limited environmental resources or carrying-capacity environment, and are more stable and permanent members of the community (9).

Some postharvest rots result from preharvest latent infections, especially in tropical and subtropical regions where environmental conditions in the field are particularly conducive to fruit infection. Controlling rots resulting from preharvest latent infections with postharvest treatments is difficult. Nevertheless, successful control of latent infections by postharvest applications has been reported (103, 106, 111). *Bacillus* spp., which are dominant colonizers of the phylloplane and fructoplane, can control latent infections by *Colletotrichum* spp. on mango and avocado (43, 44, 108). Rapid colonization by the antagonist may not be necessary for the control of rot resulting from latent infections. In the biocontrol of anthracnose of avocado, *Bacillus subtilis*, applied as a dip or a wax formulation removes nutrients from the immediate surrounding of the appressorium of *Colletotrichum gleosporioides* to ensure maintenance of the dormancy, as demonstrated by Korsten et al. (111).

The main strategy used to suppress postharvest fruit decay is the postharvest application of antagonists to prevent pathogens from infecting fruit wounds after harvest, but postharvest decay can also be suppressed by field application of biocontrol agents (16, 110, 114, 173). Field applications have been successful against

anthracnose caused by *Colletotrichum* sp. on mango and avocado (108, 111), and to a lesser extent against *Pezicula malicorticis* on apples (114), using multiple applications of the antagonist in the orchard. Although sometimes useful in control postharvest decays from wound infection (16, 172), preharvest applications of antagonists may serve only as a supplement to their postharvest application. To control postharvest decays originating from latent infection in the orchard, the application of antagonists must be repeated. Korsten (107) showed that three sprays with *B. subtilis* controlled anthracnose of avocado as well as three copper sprays, but the volume of biocontrol product required to spray the trees was large. In addition, continuous application of *B. subtilis* over a four-year period resulted in a gradual build-up of the antagonist, reduction of the level of pathogen inoculum, and more effective control.

## PATHOGENS VULNERABLE TO BIOCONTROL

The most successful biocontrol systems against plant pathogens, *Agrobacterium radiobacter* strain K-84 (currently strain K-1026) against crown gall caused by *A. tumefaciens*, and *Phlebia gigantea* against annosus root rot of pine, are based on a single application of the biocontrol agents directly to the wounds (uncolonized in the case of *Phlebia*), the site of pathogen entry (105, 145). Wounds are "an ideal environment for antagonists because they provide a moist, nutrient-rich substrate with initially no competition from other microorganisms" (18). However, to sustain such a dominant position in wounds requires a pre-emptive colonization strategy by the antagonist to ensure that the environment is "managed" for as long as possible so that it remains unfavorable for other microorganisms. Natural antagonists can be classified with the *r-K* gradient of population strategies within microbial communities. The properties of an effective biocontrol agent will therefore depend on the setting in which it is intended to function (14). The ability to rapidly colonize can be ascribed to the typical *r* strategist. *Pseudomonas* spp. and many yeasts are typical *r*-strategists, which grow rapidly, dominate, and colonize the new niche where resources are temporarily abundant. Biological agents that are *r*-strategists can be compared to a protectant fungicide that is in place before onset of the pathogen infection cycle (14). Where a pathogen has already invaded the plant host, a more competitive species with a high tolerance to abiotic stress will be required.

Plant pathogens are typically spread across the *r-K* range of characteristics. Brown rot (caused by *Monilinia fructicola*) of peach, which can originate from both latent infections in the orchard and wounds made after harvest, would thus require both *r* and *K* strategists applied in both preharvest and postharvest settings to achieve optimal control (139, 140). Wound-invading necrotrophic fungi such as *Penicillium expansum* and *B. cinerea*, which cause postharvest blue mold and gray mold on apples and pears, respectively, require nutrients for germination and initiation of the pathogenic process. This requirement makes them suitable for biocontrol through nutrient competition. The pome fruit system is well suited for

biocontrol, and many successful attempts have been reported from various laboratories (23, 26, 30, 32, 34, 55, 59, 80, 90, 101, 118, 123, 142, 147, 150, 152, 162, 178, 191). In addition, decay was reduced significantly after preharvest orchard application of antagonists, which allowed antagonist populations to become established on fruit surfaces before anticipated wounding during harvest (16, 172). A biocontrol system for reducing decays of citrus is also promising (6, 19, 27, 47, 48, 75, 125, 153, 155, 184), especially in locations where most decays are caused by wound-invading *Penicillium italicum* and *P. digitatum*, which cause blue mold and green mold, respectively. However, tests in citrus packinghouses indicated that biocontrol alone cannot provide adequate control and must be combined with diluted fungicides or other methods of control (47, 48). Other successful attempts have been made to control wound-originating postharvest decays on cherry (31, 179; L. Grant, EcoScience Corp., unpublished information), grape (15), tomato (28), kiwi fruit (33), strawberry (72, 78), and banana (42, 103).

The potential for biological control of postharvest diseases of vegetables has been reviewed recently (91).

## SEARCH FOR THE ANTAGONIST

### Sources and Isolation of the Antagonist

The fruit surface is an excellent source of naturally occurring antagonists against postharvest fruit decay. For example, searching for antagonists on healthy apples in the orchard and storage, "a place where a disease can be expected but it does not occur" (10), resulted in the isolation of many ecologically fit bacterial and yeast antagonists effective against postharvest decays of apple (80, 83). Similar isolations from other fruits also yielded effective antagonists against postharvest fruit decay pathogens (3, 6, 19, 72, 75, 104, 118, 141, 175, 184, 195). Isolation of the antagonists can be improved by using fruit from unmanaged orchards (59, 85), where natural populations have not been disturbed by chemical usage, and the pool of potential antagonists is greater than in a chemically managed orchard (160). A variety of enrichment procedures have been used that favor isolation of microorganisms growing efficiently on the substrate, which occurs at the infection site (wound) that must be protected. These include isolation from natural cracks on the fruit surface (W.J. Janisiewicz, unpublished); agar plates containing apple juice that were seeded with fruit washings (83); fruit wounds treated with fruit washings and incubated for several days (187); freshly made wounds on apples in the orchard that were exposed to colonization by fruit-associated microbiota from one to four weeks before harvest (85); and from an apple juice culture resulting from seeding diluted apple juice with the orchard-colonized wounds and repeated reinoculation to fresh apple juice.

The fructoplane has provided the most abundant and most desirable source for isolating antagonists against postharvest fruit pathogens. However, the antagonists may also come from other closely related or unrelated sources. The phylloplane

has also been a good source of antagonists, as it may share part of the resident microflora of fruits as well as contain other microorganisms dislodged from the fruit (59, 80, 109, 110, 163, 196). Screening collections of yeast (64) or starter cultures used in the food industry (138) may also yield effective antagonists. Soil also may be an abundant and diverse source of antagonists. *Bacillus subtilis* B3, a contaminant found during routine isolation of fungi from peach roots and identified for its inhibition of *M. fructicola* on agar plates (J. Barrat, personal communication), is a very effective strain for biocontrol of brown rot of peach (139). B3 was the first strain studied extensively for biocontrol of postharvest diseases, and early successes in biocontrol of brown rot on peach with this strain (139) stimulated interest in biocontrol of postharvest diseases of fruits in general.

## Selection Criteria

Cook stated, "a serious shortcoming of microbial biocontrol as currently tested and studied is that too few strains are examined" (36). To some extent, this shortcoming is also true for BCPD. Although screening of large numbers of organisms has been reported, only recently has more effort been devoted to testing the biocontrol potential of various strains of the same species of yeast and bacterial antagonists (95, 149).

For biocontrol to be successful, the conditions that favor a potential antagonist should be the same or similar to those that favor the pathogen. The best antagonists perform well over the full spectrum of conditions conducive to pathogen development. Strains of an antagonist species can be compared for effectiveness in controlling fruit decay, and for phenotypic characteristics that are useful in determining their commercial potential. For example, differentiation criteria for decay control on apple can include the biological control efficacy of the strains, spectrum of activity (pathogens to be tested, cultivar range, fruit maturity stages), ability to colonize wounded and sound fruit surfaces under various conditions, utilization of substrates occurring in fruits, or growth at cold storage temperatures and at 37°C (human body temperature). Biocontrol agents are more acceptable if they can be applied together with current practices, and information on the compatibility of the biocontrol agents with chemicals used in the postharvest system, e.g., antioxidant diphenylamine (DPA) used for control of superficial scald of apple, a physiological disorder, or flotation salts used to increase buoyancy of pears during handling in water, should be developed. Additional criteria may include resistance to environmental stress in the orchard application of biocontrol agents (16, 77, 114, 172, 174), and pathogenicity of the antagonists to fruits, since strains of some antagonists with good biocontrol potential, e.g., *Aureobasidium pullulans*, can cause minor decay and russetting on some fruits (51, 124, 146).

Since the antagonists are applied to consumable products (produce), they must meet strict requirements for human safety. The initial determination of the antagonist's safety is based on the available literature and should be made early in the program, before committing time and resources into further research. The final

determination is made during commercial development of the product, and is based on an elaborate and costly toxicological profile.

Genetic and physiological niche requirements can vary among strains within a species, and intraspecies variation is poorly understood. Some species are isolated repeatedly from the same type of fruit at various geographical locations. For example, of 13 yeast species reported to be residents of apple (40), 7 (30, 32, 59, 76, 93, 95, 114, 118, 127, 147, 192) were isolated and reported to have strong antagonistic activity against postharvest decays of pome fruits by laboratories at different geographical locations. The biocontrol potential of strains of the same species from different locations may well vary, and this subject warrants in-depth investigation. In fact, the biocontrol potential and physiological and genetic diversity of the yeast *Metchnikowia pulcherrima* appears to be great even at a single geographical location, and provides a wide variety of organisms with desirable biocontrol traits (95).

Thus, in developing biocontrol, the key requirements for successful commercialization of an antagonist must be well defined, and strain searches should continue until adequate strains are found that meet all requirements. Some of those requirements are intuitive (186), but others involving mass production, formulation, application, and distribution require a more intimate knowledge of commercialization of microbial pesticides (66, 102, 136, 166). In this regard, a significant benefit can be derived from developing, early in a program, a liaison between a scientist working on developing a biocontrol system and the industry that can commercialize the biocontrol agent (166).

## MECHANISMS OF BIOCONTROL

The mechanisms of BCPD are poorly understood, mainly because relatively few attempts have been made and appropriate methods to study microbial interactions in wounds of fruit are lacking. Various mechanisms have been described, including antibiosis, production of lytic enzymes, parasitism, induced resistance, and competition for limiting nutrients and space. Often, more than one mechanism was implicated, but in no case has a sole mechanism been found responsible for biological control.

The antibiotics iturin, produced by *B. subtilis* B-3 (70), and pyrrolnitrin, produced by *Pseudomonas cepacia* LT-4-12W (94), reduced in vitro growth and conidia germination of the stone fruit pathogen *M. fructicola*, and pome fruit pathogens *P. expansum* and *B. cinerea*, respectively. Both strains controlled fruit decays caused by the respective pathogens (94, 139), and strain LT-4-12W also controlled various decays on citrus (155) and stone fruits (156). These fruit decays were also controlled by applications of the respective antibiotics alone (79, 138). However, the significance of the antibiotics in these biocontrol situations is not clear, since strain LT-4-12W still provided substantial control of blue mold decay on oranges inoculated with laboratory-derived mutants of *P. italicum* resistant to pyrrolnitrin (J.L. Smilanick, personal communication). No analogous tests were conducted with iturin. The mechanism(s) of biocontrol of *Pseudomonas syringae* strains

ESC-10 and ESC-11 (formerly known as L-59-66) used in BioSave products has not been elucidated. Bull et al. (22) showed that on some media both strains can produce syringomycin E, which is inhibitory to a variety of fungi, and that the purified compound can control green mold of lemons. However, the role of syringomycin E in biocontrol is in doubt because efforts to isolate this compound from fruit wounds treated with the antagonist have been unsuccessful (C.T. Bull, personal communication), and rapid growth and colonization of the wounds was important for biocontrol. This suggests that competition for nutrients and space may have played a major role (21).

The attachment of antagonists to pathogen hyphae has been suggested as an important factor in competition for nutrients between the antagonist *Enterobacter cloacae* and *Rhizopus stolonifer* on peach (193), and between the antagonistic yeast *Pichia guilliermondii* and *P. italicum* on citrus fruit (8), or in the lysis of *B. cinerea* by  $\beta$ -1,3 glucanase produced by *P. guilliermondii* on apple (190). Given the very high density ( $10^{10}$  cfu/ml) of *E. cloacae* needed to inhibit conidia germination and decay, exclusion may also play an important role in biocontrol. The production of ammonia by this antagonist has been reported as a mechanism of biocontrol in another system (65) and should be given consideration in the fruit system as well. To determine the significance of attachment and  $\beta$ -1,3 glucanase in biocontrol by *P. guilliermondii*, confirmatory tests in fruit wounds are needed. Filonow (62, 63) showed that the apple volatile, butyl acetate, stimulated adhesion to membrane filters and germination of conidia of *B. cinerea* and increased apple decay. The antagonists *Cryptococcus laurentii* and *Sporobolomyces roseus* but not the baker's yeast *Saccharomyces cerevisiae* used butyl acetate as a food source and reduced these stimulatory effects in vitro. These effects, however, have not been shown in apple wounds owing to technical difficulties in conducting this type of experiment. In addition, studies with various sugars occurring in apple showed greater uptake and utilization of  $^{14}\text{C}$  fructose, and stronger inhibition of conidia germination in diluted apple juice by the antagonists other than *Saccharomyces cerevisiae*, suggesting that competition for nutrients may play an important role (61).

Jijakli & Lepoivre (100), and Graves et al. (69) have shown that the yeast *Pichia anomala* strain K, effective in the control of gray mold of apple, increased production of exo- $\beta$ -1,3-glucanase threefold in the presence of cell wall preparations of *B. cinerea* in apple wounds, reducing lesion size by more than half compared to the antagonist alone. This strengthens the hypothesis that exo- $\beta$ -1,3-glucanase is involved in the biocontrol of gray mold by this antagonist. Higher  $\beta$ -1,3-glucanase and chitinase activity was also detected in apple wounds treated with strains of an antagonist, *A. pullulans*, effective in controlling various decays on apple, table grape, and other fruits (25, 76). The increased amount of these enzymes was attributed to higher production by the antagonist and to the induction of the enzymes in the fruit itself. The source of these enzymes and their significance in biocontrol warrant further investigation. Further evidence for the significance of these enzymes in biological control could come from studies evaluating disease suppression by mutants with a disrupted  $\beta$ -1,3-glucanase gene. *A. pullulans* may also



produce antibiotic aurebasidins, whose role in biocontrol should be considered (171). *Erwinia herbicola* strains B66 and B90, which controlled blue and gray mold of apple, demonstrated taxis to the conidia and germ tubes, inhibited conidia germination, and lysed germ tubes of *B. cinerea* and *P. expansum* in diluted apple juice (20, 162). These interactions were not as apparent in undiluted apple juice, suggesting that competition for nutrients may be important in this system. The yeast *Candida famata* reduced green mold decay (caused by *P. digitatum*) on oranges and increased the phytoalexins scoparone and scopoletin 12-fold in fruit wounds after four days when inoculated alone (7). However, the significance of phytoalexins in this biocontrol is not clear because of their relatively slow production. Electron microscopic observations indicate rapid colonization and partial lysis of the pathogen's hyphae by the antagonist.

In only a few of these examples has work continued beyond the initial reports to fully explain the biocontrol mechanisms, mainly because developed methods to study mechanisms of biocontrol in BCPD, in particular competition for nutrients, are lacking. In almost all cases, nutrient competition was reported to play a significant role, but it is difficult to separate from other mechanisms. However, recent reports on developing biological sensors (121) and using natural substrates in in vitro cylinder-well tests for studying antagonist-pathogen interactions (96) may be useful in studies evaluating microbial competition for nutrients in fruit. A biological sensor, composed of a nutrient-responsive promoter fused to a reporter gene, could be used to assess the spatial distribution and availability of nutrients in fruit wounds at a critical time for pathogen. Reporter genes encoding the Green Fluorescent Protein (GFP) or ice nucleation protein (132) are especially useful for studies evaluating gene expression by bacterial antagonists on and in plant tissues. In cylinder-well tests, the antagonist and pathogen are separated by a membrane and immersed in fruit juice, which can flow through the membrane. After an incubation period, the pathogen can be removed and evaluated for viability, ability to infect fruit, or susceptibility to other mechanisms of biocontrol.

Applied microbial ecology was an initial driving force in the commercial development of BCPD of fruits, but further progress in improving biocontrol will largely depend on the basic understanding of the mechanism(s) of biocontrol.

## ENHANCEMENT OF BIOCONTROL

In comparison to the field environment, post-harvest environments are well defined; abiotic and biotic factors can be determined with relative ease and manipulated to an antagonist's advantage. Although the mechanism(s) of biocontrol have not yet been fully explained and, to date, there have been only a few attempts to exploit these mechanisms to improve postharvest biocontrol (23, 97), reports on the mechanism of BCPD suggest that competition for nutrients and space plays a major role in most cases (23, 31, 46, 61, 64, 85, 96, 110, 181, 190). Rapid colonization of fruit wounds by the antagonists is critical for decay control, and manipulations

leading to improved colonization enhance biocontrol (97, 130). Within microbial communities, interactions are density dependent, and more than one type of interaction can occur at any one time, depending on the growth phase of different microorganisms, population density, and species diversity. Three different types of interactions, competition for nutrients, competition for space, and inhibition by secondary metabolites, were observed with preharvest sprays of *B. subtilis* to control *C. gleosporioides* on avocado (110). The main approaches used to improve BCPD are (a) manipulation of the environment, (b) use of mixed cultures of antagonists, (c) physiological and genetic manipulation of antagonists, (d) combining field and postharvest applications, (e) manipulation of formulations, and (f) integration with other methods.

## Manipulation of the Environment

It is possible to manipulate the physical and chemical environment to the advantage of antagonists in storage, but fruit quality must be maintained. Temperature, humidity, and often gas compositions are predetermined to maintain fruit quality, and the antagonists must be well adapted to these conditions (44, 178). Fruits are often treated and/or handled in a water suspension before, during, and after storage, which provides an excellent opportunity to modify the environment. Nitrogen is likely to be a limiting nutrient in the carbon-rich environment of apple and pear wounds. The addition of L-asparagine and L-proline enhanced populations of the antagonist *P. syringae* in wounds of mature apple fruit more than tenfold during the critical first 24 h at room temperature and the first month of storage at 1°C, resulting in reduction of blue mold decay from as much as 50% to 0% (97). These nutrients were selected after in vitro screening of various nitrogenous compounds for preferential stimulation of the antagonist's growth over mycelial growth and conidial germination of the pathogen *P. expansum*. The addition of a nutrient analog 2-deoxy-D-glucose, which is taken up by the pathogen but is not metabolized, thus inhibiting pathogen growth and giving advantage to antagonists *P. syringae*, *S. roseus*, and *Candida saitoana*, improved biocontrol of decays of apple and citrus, respectively (58, 84). The addition of siderophores may reduce apple decay by sequestering iron required for germination of some postharvest pathogens (23). It may also stimulate production of the antagonist's siderophores by creating an iron-deficient environment at the wound site. Despite its great potential for improving biocontrol, manipulation of the chemical environment has not been widely exploited. This may be attributable to limited knowledge of the mechanisms of biocontrol. If nutrients limiting growth of pathogens or antagonists were known, then limiting nutrients or substances could be manipulated to stimulate antagonist populations and/or biocontrol mechanisms.

## Applying Mixed Cultures

It has been difficult to select individual strains with a broad spectrum of activity against major and minor pathogens that are effective when used on fruits at various

maturity levels. Mixtures of compatible strains may be needed to provide the necessary spectrum of activity. Components of the microbial community that contain the desired antagonistic attributes might be reconstructed by selecting them from fruit wounds, growing them in culture, and applying them to fruit surfaces as a mixture. In this case, the antagonistic action will result not from an activity of one species but rather from the action of a community of microorganisms that suppress a target pathogen through different mechanisms of action. However, such community reconstruction may be very challenging, as some microorganisms are incapable of independent growth under common cultural conditions (5). A better understanding of microbial communities on fruit is needed to take full advantage of this resource.

The application of antagonist mixtures has reduced variability and improved efficacy of biocontrol in many systems, some of which include pathogens that infect fruit (67, 71, 120, 135, 144, 151). With the exception of work on biocontrol of anthurium blight, where microbial communities of guttation fluids were screened for biocontrol activity (68), no special criterion was used to select antagonists for mixtures. On apple, a broader spectrum of pathogens was controlled (81), and less total biomass of the antagonist was needed to control decay when antagonists were applied in mixtures instead of individually (85, 87, 114). The mixtures were composed of antagonists paired at random (81, 114), or after screening for minimum mutual niche overlap in the utilization of nutrients, assessed on BIOLOG standard plates (85) or on customized BIOLOG plates containing nutrients occurring at the wound site (W.J. Janisiewicz, unpublished data). The BIOLOG approach reduced selection of strains competing for the same nutrients, but did not eliminate potential negative interactions that may result from the production of secondary metabolites (114). Thus, to determine further compatibility of the strains selected for a mixture, it is important to conduct coexistence studies using the De Wit displacement series (189) in fruit wounds (85). Antagonists selected for mixtures may also be obtained from microbial succession at the wound site. First, the sequence in colonizing freshly made apple fruit wounds in the orchard just before harvest must be determined and then the organisms for the mixtures selected from the succeeding organisms. A large number of such organisms were isolated and screened for antagonistic activity directly on apple fruit. The largest number of organisms antagonistic to *P. expansum* were found in wounds sampled closest to harvest (85). The isolated antagonists were ecologically suited to the chemical environment of the fruit, and selection of strains suited for the stable physical environment in storage was not difficult. We can also explore relevant examples of microbial interaction in food products. For example, the succession of yeasts in the apple cider-making process may closely resemble those at the wound site where apple juice is the main substrate (13, 17). After exhausting limiting nutrient(s) by one organism, another organism, originally less competitive but not requiring or able to synthesize the limiting nutrient, may take over colonization of the wound, further depleting nutrients utilized by the pathogen. To predict competitive interactions between antagonists in a mixture and between a

pathogen and the antagonists, the nutrient composition at the wound site must be known.

The benefits of using an antagonist mixture are clear, but implementation of this approach requires approval from the industry producing biocontrol agents, because it entails a doubling of the cost to commercialize the antagonist mixture as compared to a single antagonist. The economic viability of this approach is favored if mixtures include at least one antagonist that has already been commercialized (W.J. Janisiewicz, unpublished findings).

## Physiological and Genetic Manipulations

Physiological manipulation of antagonists has been focused on improving their ecological fitness, which is particularly important in orchard applications where environmental conditions fluctuate widely. Teixidó et al. (173) reported that populations of low water activity ( $a_w$ ) tolerant cells of the antagonist *Candida sake* obtained from growth on media modified for low ( $a_w$ ) and applied to apples 2 days prior to harvest, increased until harvest time, whereas those from nonmodified media remained relatively unchanged. Population increases of both types of cells were similar during four months of cold storage of the fruit. Although no significant difference in biocontrol of blue mold on apples was observed between the two types of cells, this study showed the potential of using this approach to increase antagonist populations, which could enhance biocontrol in other systems where antagonist populations declined after orchard application (16).

Physiological manipulation may also be used to enhance mechanisms of biocontrol. This has been demonstrated for soil antagonists, where the addition of zinc increased production of the antibiotic, phenazine, which is a mechanism by which *Pseudomonas fluorescens* suppresses take-all of wheat caused by *Gaeumannomyces graminis* var. *tritici* (154). In a postharvest system, Calvente et al. (23) reported that iron sequestration by a siderophore, rhodotorulic acid, produced by the yeast *Rhodotorula glutinis*, is responsible for biocontrol of *P. expansum* on apples. The addition of the siderophore to the antagonist suspension further increased biocontrol of blue mold on apples, whereas the addition of iron reduced biocontrol. This study demonstrates that competition for iron can be an important biocontrol mechanism on fruit.

Genetic manipulation of antagonists to improve BCPD is a field in its infancy. Current efforts are focused on developing efficient transformation procedures for bacterial and yeast antagonists and inserting genes for tracking the antagonist in the environment rather than enhancing biocontrol (12, 133, 194). However, attempts to overexpress genes involved in biocontrol, e.g., lytic enzymes, or engineering strains with desired biocontrol traits (M. Wisniewski & W.J. Janisiewicz, unpublished results), may soon yield positive results.

## Preharvest Applications

There are two distinct approaches in applying biocontrol agents in the field to control postharvest decays of fruits. In one, antagonists developed for postharvest

application are applied just before harvest. The intent is to precolonize the fruit surface with an antagonist immediately before harvest, so that wounds inflicted during harvest can be colonized by the antagonist prior to colonization by a pathogen (77). In the other approach, antagonists, selected for field application, are applied throughout most of the fruit development, to reduce latent infections that can originate as early as bloom time and cause fruit decay after harvest when the natural mechanisms of resistance have broken down. These two approaches are different from biological control of fruit decays in the field, which are beyond the scope of this review (50). In both approaches, many of the advantages of postharvest application of biocontrol agents discussed earlier are lost. However, both approaches have had some successes. The antagonistic yeasts *Cryptococcus infirmo-miniatus*, *C. laurentii*, and *R. glutinis*, applied to d'Anjou and Bosc pears in the field 3 weeks before harvest maintained high population densities through harvest and, in the best cases, reduced gray mold on Bosc pear from 13% to as little as 4%, and on d'Anjou pear from 7% to as little as 1% on fruit that were wounded after harvest and stored for 4 months at 0.5°C (16). Given such a high reduction in decay from just one preharvest application of the antagonist, it would be interesting to determine how much more protection can be achieved by an additional application after harvest, when postharvest fungicides are routinely applied. Populations of antagonists such as *A. pullulans*, *R. glutinis*, and *B. subtilis* adapted to variable environmental conditions, increased after application to apples in the orchard, and were maintained at relatively high densities on fruit in cold storage, except for *B. subtilis*, which declined (114). Mixtures of these antagonists controlled blue and gray molds and bull's-eye rot caused by *P. malicorticis* as effectively as the fungicide Euparen, and were more effective than the individual antagonists in tests on apples after harvest. Applying mixtures of these antagonists in the orchard may be useful for controlling postharvest decays, especially if applied early in the growing season, which may reduce latent infection caused by *Pezicula* spp. *C. sake* CPA-1 reduced blue mold by ~50% on wounded apples inoculated with the antagonist 2 days before harvest and then inoculated with *P. expansum* before placing in cold storage for 4 months (172). Although these procedures would be unlikely under commercial conditions, this work confirmed that wounds precolonized by the antagonist in the field are much more difficult to infect by the pathogen. Here too, it will be interesting to see how much more decay control can be achieved with an additional antagonist application after harvest.

The control of postharvest decays of strawberries has been very difficult, even with preharvest fungicidal applications. Field infections are considered to be the main source of fruit decay because *Botrytis* infections can occur from bloom to harvest and often develop into decay on mature fruits (98). Field applications of various antagonists including *Gliocladium roseum* (169), *Trichoderma hazianum* (113, 176), *B. subtilis*, and *B. licheniformis* (L. Korsten, unpublished data), yeast, and other bacteria (F. Takeda & W.J. Janisiewicz, unpublished data) from bloom until harvest have had variable success. Less variable results were obtained in controlling fruit decays in greenhouse strawberry culture, and antagonists on flowers and fruit remained at higher and more stable populations under controlled

greenhouse conditions (119; F. Takeda & W.J. Janisiewicz, unpublished data). The best control, however, was obtained with the application of pyrrolnitrin, a secondary metabolite from the biocontrol agent *P. cepacia*, on harvested fruit to control postharvest decays (170). Anecdotal evidence suggests that postharvest infection originating from wounds made by pickers, basket abrasions, and handling may be a major cause of strawberry decay (41; W.J. Janisiewicz, personal observation). Perhaps biocontrol efforts should be focused on the control of these kinds of infections. In this respect, there is a recent, promising report on the successful biocontrol of strawberry decay with postharvest application of *Candida reukaufii*, *C. pulcherrima*, and strains of Enterobacteriaceae isolated from strawberry fruits (72).

In the examples above, significant biocontrol of postharvest decays on the same kind of fruit was achieved with both field and postharvest applications. Combining field and postharvest application of biocontrol agents should lead to even more effective control of postharvest decays.

## Manipulation of Formulations

Formulations have a profound effect on biocontrol agents and products, including shelf life, ability to grow and survive after application, effectiveness in disease control, ease of operation and application, and cost (66). Formulation of microorganisms for biocontrol of plant pathogens is undeveloped compared with other applications of microorganisms. Formulations of commercial products for postharvest applications remain mostly proprietary, but research in the public sector is recording progress on formulations for improving on the viability, efficacy, and shelf life of biocontrol agents. Certain freeze-drying protective agents and rehydration media enhanced the viability of the antagonist *Pantoea agglomerans* strain CPA-2, effective against blue mold and gray mold of pome fruits (38). The greatest protection from freeze-drying injury to the antagonist was provided by 5% trehalose, with the survival of over 60% of the cells, whereas 10% skim milk was the best rehydration medium, resulting in a 100% recovery of the freeze-dried bacterium. At high initial cell densities of the antagonist ( $10^{10}$  cfu/ml), sucrose was a very effective protectant, whereas rehydration media had no effect on recovery. Survival of cells of the antagonistic yeast *C. sake* was improved from 0.2% to 30–40%, by using freeze-drying protective media consisting of skim milk and other protectants, such as 10% lactose or glucose, and 10% fructose or sucrose. However, the shelf life of the product has been poor and needs additional improvement (1). The addition of xanthan gum to *A. pullulans* L47, applied to strawberries in the field from bloom to fruit at the green stage, improved survival of the antagonist and increased biocontrol of storage rot caused by *B. cinerea* (78).

Formulations can influence the survival and activity of biocontrol agents on fruit surfaces and in wounds. In some spore-forming bacteria such as *Bacillus* species,

breaking the endogenous dormancy of the formulated product may be a factor in the speed of action, which is crucial in postharvest biocontrol. Formulations that include wetters (humectants) to facilitate reabsorption of moisture from air may reduce this problem (102). Wetters not only make water spray stay on plants but, like oil carriers, they also enable organisms to reach otherwise inaccessible places such as depressions, stomata, and lenticels, thereby improving the chances of establishing antagonists for disease control. Oil carriers are expensive, but formulations containing oils can enhance the reliability of biological control agents (102). Ultra low volume (ULV) sprays give good cover in foliage canopies under suitable conditions and are relatively cost effective. Also worth exploration is the use of natural pigments such as melanins, which are nature's super sunscreen and also improve dessication tolerance and protect against hydrolytic enzymes. Research is needed to determine the value of each additive alone and also in the presence of other ingredients.

Improvements in formulations can result in substantial benefits to biological control, but research in this area has been limited. A more systematic approach to improve formulations is required and could be undertaken with knowledge of the biocontrol agent's mode of action and the windows for effective use, especially under field conditions. An understanding of agricultural use and market factors is also needed to improve biocontrol formulations. A shelf life of less than 6 months requires direct order service, whereas 0.5–2 years is adequate for conventional off-shelf sales. The biocontrol products BioSave100 and 110 have been formulated as frozen pellets and as a wettable powder. The frozen formulations require a cold chain and extensive field services to assure high quality of the product (L. Grant, EcoScience Corp., personal communication). The shelf-life limits of each biocontrol agent is probably controlled by its genetic composition. These limits might be extended both by evaluating more strains and by genetic engineering with genes affecting survival traits.

## Integration with Other Methods

During the past few decades, many attempts have been made to develop non-fungicidal methods to control postharvest decays on various commodities. They include environmental modification such as storing commodities at temperatures suppressive to pathogen development, modifying relative humidity and the atmosphere, and treatment with hot air or water (60, 161, 164); inducing resistance by applying elicitors (54) or UV irradiation (45, 168); applying substances generally regarded as safe (GRAS) (35, 157–159); and sterilizing fruits and handling water with UV irradiation (134) or ozone (177). However, none of these methods, when used alone, provided satisfactory levels of decay control, although some appeared to be very useful when applied in combination with biological control, resulting in additive or even synergistic levels of decay control.

Applying a 2% solution of calcium chloride together with the yeast antagonist *Candida* spp. enhanced biocontrol of gray and blue molds on apples (128, 191), but calcium chloride solution alone did not reduce decays. However, application of 68 mM  $\text{CaCl}_2$  to grapefruit reduced the incidence of green mold decay by 43%, and in combination with an antagonist *P. guilliermondii* strain US-7 by 97% (49). Pressure infiltration of 0.27 M calcium chloride into apples increased calcium content of the mesocarp threefold and reduced blue mold decay on Golden Delicious apples inoculated with *P. expansum* after 6 months in cold storage by as much as 50% (88). Combining calcium chloride pressure infiltration with the application of the antagonist *P. syringae* (isolate ESC-11 used in BioSave110) resulted in greater control than the individual treatments (88). Integrating these two treatments has the added benefit of increased Ca to alleviate physiological storage maladies such as bitter pit, and reduced amounts of both products to be used without compromising decay control.

Prestorage hot air treatment (38°C for 4 days) of apples reduced or eliminated blue mold decay caused by *P. expansum* and gray mold decay (60). Heat had eradicated activity on decay of apples inoculated with *P. expansum* 12 h before heating. Heat also improved biocontrol with heat-tolerant yeasts when applied to apples up to 24 h after inoculation with the pathogen (116). Combining the heat treatment with Ca infiltration and then applying the biocontrol agent *P. syringae* ESC-11 was most effective in reducing blue mold decay compared with individual or other treatment combinations, when mature fruit, after 6 months in storage at 1°C, were inoculated with *P. expansum* alone or in a mixture with the antagonist. The heat treatment provided little residual protection, but the residual protection provided by Ca and the antagonist added to the control by heat (117). When antagonists were applied to apple wounds before heat treatment, the heat reduced populations of *P. syringae* and increased populations of the two heat-tolerant yeasts more than tenfold (117). After removal from the heat and placement of fruits in cold storage, populations of *P. syringae* increased, but not to the original application level, and those of the yeasts continued to increase. The addition of the heat treatment improved control of blue mold for this and two other heat-tolerant yeast antagonists. Hot water (45–56°C) dip treatments are used commercially in citrus and mango packinghouses in South Africa to reduce postharvest decays. The efficacy of control could be enhanced by the addition of *Bacillus* spp. to the hot water dips (L. Korsten, unpublished data).

Chitosan and its derivatives, including glycolchitosan, were reported to inhibit fungal growth and to induce host-defense responses in plants and harvested commodities (4, 185). Combining 0.2% glycolchitosan with the antagonist *C. saitoana* was more effective in controlling green mold of oranges and lemons, caused by *P. digitatum*, and gray and blue molds of apples than either treatment alone (56, 57). Pretreatment of lemons with sodium bicarbonate further increased control of green mold on the light-green and yellow lemons (57). Other fruit coatings may also be useful for further reducing decay when applied with biocontrol agents (11, 39). For example, a fruit coating containing sodium salts of carboxy-methyl-cellulose,



sucrose esters of fatty acids, and mixed sucro-glycerides and soap, commercialized under the name TAL Pro-long, reduced the spread of a range of postharvest decays of pome fruits (11). The mechanism of action of TAL-Pro-long is not fully understood, but it reduces ripening and extends the natural resistance to invasion by the pathogen in storage.

Irradiation of root vegetables such as carrots (129) or sweet potatoes (167) with UV-C (wavelength below 280 nm) induced resistance and reduced postharvest decays. Attempts to irradiate pome, stone, and citrus fruit with UV-C and combine irradiation with biological control were largely unsuccessful (185). This may be attributed to a narrow range of the optimum UV-C irradiation dose, which is specific to different fruits, and variations of the resistance response with fruit maturity and storage temperature (45). The physiology of root tissue is very different from that of fruit, with the response of roots to UV-C irradiation more consistent (129, 167). This suggests that combining UV-C and biocontrol treatments may have a greater effect in controlling postharvest decays on roots than on fruits, and the role of the UV-C on fruits will be restricted mainly to its phytosanitary effect of reducing the survival of pathogen propagules.

GRAS substances such as sodium carbonate, sodium bicarbonate, and ethanol reduced conidial germination of *P. digitatum*, the causal agent of green mold of citrus (157, 158). Combining treatments of 3% sodium carbonate and the antagonist *P. syringae* ESC-10 was superior to either treatment alone in controlling green mold on citrus (159). This combination overcomes the significant shortcomings of both individual treatments. The antagonist alone is a poor eradicator and is usually incapable of controlling green mold on fruit inoculated with the pathogen 24 h before treatment with the antagonist. In contrast the carbonate salts control these infections (157). Carbonate salts, on the other hand, do not provide persistent protection from reinfection after treatment, whereas the antagonist persists for long periods after application and protects fruit from reinfection. Ethanol at 10%, in combination with ethanol-resistant *S. cerevisiae* strains 1440 and 1749, isolated from wine and ensile acorns, respectively, reduced the incidence of gray mold decay on apples from more than 90% to close to 0%, whereas either treatment alone did not reduce decay (122). The same concentration of ethanol reduced green mold of lemons to less than 5% (158). It will be interesting to determine the effect of ethanol in combination with an ethanol-resistant biocontrol agent on green mold of lemon.

Some non-fungicidal methods for control of postharvest diseases have been used on a small scale for some time and should be easy to combine with biocontrol treatments. Examples include sodium carbonate treatment of lemons, the addition of  $\text{CaCl}_2$  to handling suspensions, and UV irradiation of pome fruit. Other approaches described above await implementation. Nevertheless, combinations of these treatments with biocontrol show great potential for enhancing control of postharvest decays of fruits.

Biocontrol product performance can be enhanced by applying the antagonist in a cascade similar to the hurdle technology strategy used in the food industry.

By incorporating various control steps along the packing line from receiving to packing, different combinations of products can be tested that more specifically suit individual packinghouses. For instance, by first drenching or fine-spraying fruit with a disinfectant such as chlorine, followed by hot water dip or ethanol spray, hot air drying, and finally biocontrol treatment incorporated into wax, effective control of mango, citrus, and avocado postharvest diseases can be provided. Creating a vacuum on the fruit surface and subsequently filling it with natural antagonists will not only reduce competition with other epiphytes, but will also stimulate the plant's natural defense system.

## DEVELOPING THE COMMERCIAL PRODUCT

Biological control of plant diseases in general and on fruit after harvest in particular is a niche market, with a relatively small profit potential. Thus, finding an industrial partner has been the first challenge to public-sector researchers seeking to commercialize a biocontrol strain. In the case of BioSave development, the effectiveness of the antagonist, a saprophytic strain of *P. syringae* L-59-66, in reducing blue mold and gray mold decay on apples and pears in a laboratory setting was demonstrated to EcoScience Corp. (Orlando, FL). Then, large-scale feasibility tests were conducted in cooperation with the company. In these tests, the antagonist was applied to fruit on a commercial packing line, and after 3 months of storage, the fruit was evaluated for the development of decay. The commercial setting of the test, involvement of industry in conducting those tests, and encouraging results were the key factors in obtaining a commitment to develop the antagonist for commercial use (W.J. Janisiewicz, unpublished data). EcoScience Corp. investigated the potential for registration and formulation of the antagonist before making this commitment. Mass production by fermentation and the biomass yield of *P. syringae* strain L-59-66 was determined before scale-up experiments. The following years were focused on determining the spectrum of activity (92; W.J. Janisiewicz, unpublished data), testing the formulations developed by EcoScience Corp., testing the final formulation in a pilot test (90), and the development by EcoScience of safety data for registration of the antagonist. To build confidence in the product within the fruit industry, pilot tests were conducted in commercial packinghouses (99, 166). Extensive technical support and quality control have been instrumental in the success of this product. BioSave use has been increasing steadily over the past five years; during the 2000–2001 storage season, over 34 million bushel boxes of produce were treated with BioSave.

The commercial development of Aspire by Ecogen-Israel Partnership, Ltd. focused on the biocontrol of postharvest decays of citrus, mainly blue mold and green mold caused by *P. italicum* and *P. digitatum*, respectively, which invade through wounds after harvest. The early research (29, 184) and the pilot test (48) in commercial packinghouses were conducted with a yeast biocontrol agent,

*P. guilliermondii* (originally described as *Debaromyces hansenii*) (126). Results indicated that a combination of the yeast with a tenfold diluted commercial rate of thiabendazole (TBZ) provided control equal to the full-strength fungicide. Subsequently, the focus has been changed to the yeast antagonist *Candida oleophila* (130, 191), previously identified as *C. sake* (187). Tests in commercial citrus packinghouses on oranges and grapefruit in Israel indicated that, like *P. guilliermondii*, *C. oleophila* gave satisfactory control of green and blue mold and sour rot caused by *Geotrichum candidum* only in combination with tenfold diluted TBZ (47). Perhaps a more effective antagonist could have been found if more emphasis was put on preliminary and secondary screening at the initial stage of research, and on selection criteria for antagonist designated for commercial development. In addition, changing the biocontrol agent during commercialization may have had an effect on the final product, as less extensive studies with the new antagonist were conducted. It is necessary to take into account as many aspects of biocontrol prior to commercialization as possible; hasty commitment to biocontrol agents without appropriate testing and classification should be avoided. The citrus system seems less prone to biocontrol than the pome fruit system. Combining yeasts with other control methods (188) appears to be a sensible approach at this time, although the citrus system may benefit greatly from more rigorous selection and manipulation of the antagonists to obtain a strain with enhanced biocontrol activity.

The research and commercial development of YieldPlus for biocontrol of pome fruit decays seems to follow the previously described example; however, little has been reported about that product.

The development of Avogreen followed a slightly different path, as this antagonist is applied in the field for postharvest biocontrol. Avogreen is a commercial formulation of the biocontrol agent *B. subtilis* isolated from avocado phyloplane, which has been registered in South Africa for control of *Cercospora* spot and anthracnose of avocado. Growers were encouraged to first test the product on a limited scale and integrate it with existing copper sprays. Support was provided to calculate dosage and adjust the existing spray schedule, which took into account equipment, cultivars, age, and history of the orchard disease profile. Developing different formulations was required to address the different needs of growers in terms of mixing, integrating with existing chemicals, and application method, which varied between low- and high-pressure equipment.

Marketing biological control products requires specialized knowledge of the target plant disease, the biological control agent, integrated disease control practices, production and storage systems, and microbial ecosystems. The success of implementing biocontrol or integrated disease management systems will depend largely on product knowledge and a thorough understanding of the complexity of the disease and postharvest environment. Distributing both the product and the knowledge necessary for its successful use will be the only effective way to ensure long-term market acceptance. These requirements are often overlooked in commercialization of biocontrol agents.

## FUTURE PROSPECTS

This review reported the success of BCPD under laboratory and commercial conditions. The fact that the commercial introduction of BCPD increased the level of familiarity with this method of control by packinghouse personnel, regulatory agencies, private investors, and the public is important in advancing the commercial use of biological control (37). In addition, continuous increases in the use of BioSave since its large-scale commercial introduction in 1996, without an incidence of failure, boost user confidence in commercial biological control of plant pathogens.

The success of BioSave indicates that current biological control practices can be cost effective in large packinghouse systems. However, the quantitative relationship between the populations of the antagonist and the resulting control necessitates the presence of high cell densities of the antagonist in product, thereby cutting profit margins. Furthermore, postharvest practices in the Central and Eastern United States, which differ from those used in the Pacific Northwest, result in using an antagonist suspension not as efficiently as in the Pacific Northwest. Therefore, use of BioSave may be too costly in those regions. Biocontrol must be adapted to practices in different regions.

The issue now is not if or when BCPD will be used, but how broad its use will be and how fast it will expand to different commodities. BCPD has its limitations, which may restrict its use under some circumstances, but many of those limitations may be effectively addressed; this method is amenable to manipulation, as indicated in the many examples presented above. It would be inappropriate to equate biological control with fungicidal treatment without considering the advantages and limitations of both methods, which often differ. Hastily designed experiments comparing the two methods, without giving consideration to those factors, should be avoided. Instead, focus should be on circumstances where currently developed biocontrol can be used effectively. Every effort should be made to expand its use by the various improvement methods described above, including new areas, e.g., control of foodborne pathogens (89, 115), where fungicides are ineffective.

Our current model of biological control of plant disease is based on knowledge of natural processes of the antagonist-pathogen interaction. Although this traditional model is credited with numerous successes, including BCPD, we should move to adapt aspects of biotechnology as a means to improve disease control with even safer and more effective methods. In the well-defined environment of a postharvest system, there are unique opportunities to use microorganisms as a delivery system. In the future, it may be possible to use only strains adapted to postharvest conditions and introduce genes for biocontrol activity as needed. Development of microbial strains, as in developing new cultivars adapted to our needs, may become common practice in the future (37).

The science and practice of BCPD is still in its infancy compared to fungicidal treatment or field biocontrol, but the progress made in this area during the past

decade and a half has been remarkable. If this pace continues, the use of BCPD will be greatly expanded in the future.

**The Annual Review of Phytopathology is online at <http://phyto.annualreviews.org>**

## LITERATURE CITED

1. Abadías M, Benabarre A, Teixidó N, Usall J, Viñas I. 2001. Effect of freeze drying and protectants on viability of the biocontrol yeast *Candida sake*. *Int. J. Food Microbiol.* 65:173–82
2. Abadías M, Teixidó N, Usall J, Benabarre A, Viñas I. 2001. Viability, efficacy, and storage stability of freeze-dried biocontrol agent *Candida sake* using different protective and rehydration media. *J. Food Prot.* 64:856–61
3. Adikaram NKB, Karunaratne A. 1998. Suppression of avocado anthracnose and stem-end rot pathogens by endogenous antifungal substances and a surface inhabiting *Pestalotiopsis* sp. See Ref. 101a, pp. 72–77
4. Allan CR, Hadwiger LA. 1979. The fungicidal effect of chitosan on fungi of varying cell wall composition. *Exp. Mycol.* 3:285–87
5. Andrews JH, Harris RF. 2000. The ecology and biogeography of microorganisms on plant surfaces. *Annu. Rev. Phytopathol.* 38:145–80
6. Arras G. 1993. Inhibition of postharvest fungal pathogens by *Bacillus subtilis* strains isolated from citrus fruit. *Adv. Hortic. Sci.* 7:123–27
7. Arras G. 1996. Mode of action of an isolate of *Candida famata* in biological control of *Penicillium digitatum* in orange fruits. *Postharvest Biol. Technol.* 8:191–98
8. Arras G, De Cicco V, Arru S, Lima G. 1998. Biocontrol by yeasts of blue mould of citrus fruits and the mode of action of an isolate of *Pichia guilliermondii*. *J. Hortic. Sci. Biotechnol.* 73:413–18
9. Atlas RM, Bartha R. 1998. *Microbial Ecology. Fundamentals and Applications*. Redwood City, CA: Benjamin/Cummings
10. Baker KJ, Cook RJ. 1974. *Biological Control of Plant Pathogens*. San Francisco: Freeman
11. Bancroft RD. 1995. The use of a surface coating to ameliorate the rate of spread of post-harvest fungal diseases of top fruit. *Int. Biodeterior. Biodegrad.* 36:385–405
12. Bassett CL, Janisiewicz WJ. 2001. Stability of *E. coli*/*Pseudomonas* shuttle vectors in a biocontrol strain of *Pseudomonas syringae*. *Phytopathology* 91:S7
13. Beech FW. 1993. Yeast in cider making. In *The Yeast*, ed. AH Rose, JS Harrison, pp. 169–213. London: Academic
14. Bellows TS. 1999. Foliar, flower, and fruit pathogens. In *Handbook of Biological Control: Principles and Applications of Biological Control*, ed. TS Bellows, TW Fisher, pp. 841–70. San Diego: Academic
15. Ben-Arie R, Droby S, Zutkji J, Cohen L, Weiss B, et al. 1991. Preharvest and postharvest biological control of *Rhizopus* and *Botrytis* bunch rots of table grapes with antagonistic yeasts. See Ref. 184a, pp. 100–12
16. Benbow JM, Sugar D. 1999. Fruit surface colonization and biological control of postharvest diseases of pear by preharvest yeast applications. *Plant Dis.* 83:839–44
- 16a. Bertolini P, Sijmons PC, Guerzoni ME, Serra F, eds. 1998. *COST 914—COST 915 Joint Workshop—Non Conventional Methods for the Control of Postharvest Disease and Microbial Spoilage*. Luxembourg: Publ. Eur. Community

17. Bizeau C, Drilleau JF, Michel J. 1989. *Study of interactions between yeast involved in cider-making*. Presented at Int. Symp. Yeast, 7<sup>th</sup>, pp. S169–73. Chichester, UK: Wiley
18. Blakeman JP, Fokkema NJ. 1982. Potential for biological control of plant diseases on the phylloplane. *Annu. Rev. Phytopathol.* 20:167–92
19. Borras AD, Aguilar RV. 1990. Biological control of *Penicillium digitatum* by *Trichoderma viride* on postharvest citrus fruits. *Int. J. Food Microbiol.* 11:179–84
20. Bryk H, Dyki B, Sobiczewski P. 1998. Antagonistic effect of *Erwinia herbicola* on *in vitro* spore germination and germ tube elongation of *Botrytis cinerea* and *Penicillium expansum*. *Biocontrol* 43:97–106
21. Bull CT, Stack JP, Smilanick JL. 1997. *Pseudomonas syringae* strains ESC-10 and ESC-11 survive in wounds on citrus and control green and blue molds of citrus. *Biol. Control* 8:81–88
22. Bull CT, Wadsworth ML, Sorensen KN, Tekemoto JY, Austin RK, Smilanick JL. 1998. Syringomycin E produced by biological control agents controls green mold on lemons. *Biol. Control* 12:89–95
- 22a. Burges HD, ed. 1998. *Formulation of Microbial Biopesticides*. Boston: Kluwer
23. Calvente V, Benuzzi D, de Tosetti MIS. 1999. Antagonistic action of siderophores from *Rhodotorula glutinis* upon the postharvest pathogen *Penicillium expansum*. *Int. Biodeterior. Biodegrad.* 43: 167–72
24. Cappellini RA, Ceponis MJ. 1984. Postharvest losses in fresh fruits and vegetables. See Ref. 132a, pp. 24–30
25. Castoria R, De Curtis F, Lima G, Caputo L, Pacifico S, De Cicco V. 2001. *Aureobasidium pullulans* (LS-30) an antagonist of postharvest pathogens of fruits: study on its modes of action. *Postharvest Biol. Technol.* 22:7–17
26. Castoria R, De Curtis F, Lima G, De Cicco V. 1997.  $\beta$ -1,3-glucanase activity of two saprophytic yeasts and possible mode of action as biocontrol agents against postharvest diseases. *Postharvest Biol. Technol.* 12:293–300
27. Chalutz E, Cohen L, Weiss B, Wilson CL. 1988. Biocontrol of postharvest diseases of citrus fruit by microbial antagonists. *Int. Citrus Conf.*, 6<sup>th</sup>, ed. R Goren, K Mendel, pp. 1–4. Weikersheim, Germany: Margraf Sci.
28. Chalutz E, Droby S, Cohen L, Weiss B, Barkai-Golan R, et al. 1991. Biological control of *Botrytis*, *Rhizopus*, and *Alternaria* rots of tomato fruit by *Pichia guilliermondii*. See Ref. 184a, pp. 71–85
29. Chalutz E, Wilson CL. 1990. Postharvest biocontrol of green and blue mold and sour rot of citrus fruit by *Debryomyces hansenii*. *Plant Dis.* 74:134–37
30. Chand-Goyal T, Spotts RA. 1996. Control of postharvest pear diseases using natural saprophytic yeast colonists and their combination with a low dosage of thiabendazole. *Postharvest Biol. Technol.* 7:51–64
31. Chand-Goyal T, Spotts RA. 1996. Postharvest biological control of blue mold of apple and brown rot of sweet cherry by natural saprophytic yeast alone or in combination with low doses of fungicides. *Biol. Control* 6:253–59
32. Chand-Goyal T, Spotts RA. 1997. Biological control of postharvest diseases of apple and pear under semi-commercial and commercial conditions using three saprophytic yeasts. *Biol. Control* 10: 199–206
33. Cheah LH, Hill R, Hunt AW. 1992. Potential for biological control of botrytis stem-end rot of kiwifruit with *Trichoderma* spp. *Proc. NZ Plant Prot. Conf.*, 45<sup>th</sup>, pp. 193–96
34. Combrink CJ, De Kock SL, Benic LM. 1998. Control of post-harvest decay in fruit. *Deciduous Fruit Grower* 48:1–6
35. Conway WS. 1982. Effect of postharvest calcium treatment on decay of Delicious apples. *Plant Dis.* 66:402–3

36. Cook RJ. 1992. A customized approach to biological control of wheat root diseases. See Ref. 175a, pp. 211–22, 230
37. Cook RJ. 1993. Making greater use of introduced microorganisms for biological control of plant pathogens. *Annu. Rev. Phytopathol.* 31:53–80
38. Costa E, Usall J, Teixidó N, Garcia N, Viñas I. 2000. Effect of protective agents, rehydration media and initial cell concentration on viability of *Pantoea agglomerans* strain CPA-2 subjected to freeze drying. *J. Appl. Microbiol.* 89:793–800
39. Cuppett SL. 1994. Edible coatings as carriers of food additives, fungicides and natural antagonists. In *Edible Coatings and Films to Improve Food Quality*, ed. JM Krochta, EA Baldwin, MO Nisperos-Carriedo, pp. 121–37. Lancaster/Basel: Technomic
40. Davenport RR. 1976. Distribution of yeasts and yeast-like organisms from aerial surfaces of developing apples and grapes. In *Microbiology of Aerial Plant Surfaces*, ed. CH Dickinson, TF Preece, pp. 325–59. London: Academic
41. Davis RP, Dennis C. 1979. Use of dicarboximide fungicides on strawberries and potential problems of resistance in *Botrytis cinerea*. *Proc. Br. Crop. Prot. Conf., Pests Dis.* 1:193–201
42. De Costa DM, Subasinghe SSNS. 1998. Antagonistic bacteria associated with the fruit skin of banana in controlling its postharvest diseases. *Trop. Sci.* 38:206–12
43. De Jager ES, Hall AN, Wehner FC, Korsten L. 2001. Microbial ecology of the mango phylloplane. *Microb. Ecol.* 42:201–7
44. Dock LL, Nielsen PV, Floros JD. 1998. Biological control of *Botrytis cinerea* growth on apples stored under modified atmospheres. *J. Food Prot.* 61:1661–65
45. Droby S, Chalutz E, Horev B, Cohen L, Gaba V, et al. 1993. Factors affecting UV-induced resistance in grapefruit against the green mould decay caused by *Penicillium digitatum*. *Plant Pathol.* 42:418–24
46. Droby S, Chalutz E, Wilson C, Wisniewski M. 1989. Characterization of the biocontrol activity of *Debaryomyces hansenii* in the control of *Penicillium digitatum* on grapefruit. *Can. J. Microbiol.* 35:794–800
47. Droby S, Cohen L, Daus A, Weiss B, Horev B, et al. 1998. Commercial testing of Aspire: a yeast preparation for the biological control of postharvest decay of citrus. *Biol. Control* 12:97–101
48. Droby S, Hofstein R, Wilson CL, Wisniewski M, Fridlender B, et al. 1993. Pilot testing of *Pichia guilliermondii*: a biocontrol agent of postharvest diseases of citrus fruit. *Biol. Control* 3:47–52
49. Droby S, Wisniewski ME, Cohen L, Weiss B, Toutou D, et al. 1997. Influence of  $\text{CaCl}_2$  on *Penicillium digitatum*, grapefruit peel tissue, and biocontrol activity of *Pichia guilliermondii*. *Phytopathology* 87:310–15
50. Dubos B. 1984. Biocontrol of *Botrytis cinerea* on grapevines by an antagonistic strain of *Trichoderma harzianum*. In *Current Perspectives in Microbial Ecology*, ed. MJ Klub, CA Reedy, pp. 370–73. Washington, DC: Am. Soc. Microbiol.
51. Dugan FM, Roberts RG. 1995. Etiology of preharvest colonization of Bing cherry fruit by fungi. *Phytopathology* 84:1031–36
52. Eckert JW, Ogawa JM. 1985. The chemical control of postharvest diseases: subtropical and tropical fruits. *Annu. Rev. Phytopathol.* 23:421–54
53. Eckert JW, Ogawa JM. 1988. The chemical control of postharvest diseases: deciduous fruits, berries, vegetables and root/tuber crops. *Annu. Rev. Phytopathol.* 26: 433–69
54. El-Ghaouth A. 1998. Use of elicitors to control postharvest diseases in fruits and vegetables. See Ref. 101a, pp. 131–35
55. El-Ghaouth A, Smilanick J, Brown GE, Ippolito A, Wilson CL. 2001. Control of

- decay of apple and citrus fruits in semi-commercial tests with *Candida saitoana* and 2-Deoxy-D-glucose. *Biol. Control* 20:96–101
56. El-Ghaouth A, Smilanick JL, Brown GE, Ippolito A, Wisniewski M, Wilson CL. 2000. Applications of *Candida saitoana* and glycolchitosan for the control of postharvest diseases of apple and citrus fruit under semi-commercial conditions. *Plant Dis.* 84:243–48
  57. El-Ghaouth A, Smilanick JL, Wilson CL. 2000. Enhancement of the performance of *Candida saitoana* by the addition of glycolchitosan for the control of postharvest decay of apple and citrus fruit. *Postharvest Biol. Technol.* 19:103–10
  58. El-Ghaouth A, Smilanick JL, Wisniewski M, Wilson CL. 2000. Improved control of apple and citrus fruit decay with a combination of *Candida saitoana* and 2-Deoxy-D-glucose. *Plant Dis.* 84:249–53
  59. Falconi CJ, Mendgen K. 1994. Epiphytic fungi on apple leaves and their value for control of the postharvest pathogens *Botrytis cinerea*, *Monilinia fructigena* and *Penicillium expansum*. *J. Plant Pathol.* 101:38–47
  60. Falik ES, Grinberg S, Gambourg M, Lurie S. 1995. Prestorage heat treatment reduces pathogenicity of *Penicillium expansum* in apple fruit. *Plant Pathol.* 45:92–97
  61. Filonow AB. 1998. Role of competition for sugars by yeasts in the biocontrol of gray mold of apple. *Biocontrol Sci. Technol.* 8:243–56
  62. Filonow AB. 1999. Yeasts reduce the stimulatory effect of acetate esters from apple on the germination of *Botrytis cinerea* conidia. *J. Chem. Ecol.* 25:1555–65
  63. Filonow AB. 2001. Butyl acetate and yeasts interact in adhesion and germination of *Botrytis cinerea* conidia *in vitro* and in fungal decay of Golden Delicious apple. *J. Chem. Ecol.* 27:831–44
  64. Filonow AB, Vishniac HS, Anderson JA, Janisiewicz WJ. 1996. Biological control of *Botrytis cinerea* in apple by yeasts from various habitats and their putative mechanisms of antagonism. *Biol. Control* 7:212–20
  65. Fravel D. 1988. Role of antibiosis in the biocontrol of plant diseases. *Annu. Rev. Phytopathol.* 26:75–91
  66. Fravel D, Connick WJ Jr., Lewis JA. 1998. Formulation of microorganisms to control plant diseases. See Ref. 22a, pp. 187–202
  67. Fukui R, Fukui H, Alvarez AM. 1999. Comparisons of single versus multiple bacterial species on biological control of anthurium blight. *Phytopathology* 89:366–73
  68. Fukui R, Fukui H, Alvarez AM. 1999. Suppression of bacterial blight by bacterial community isolated from the guttation fluids of anthuriums. *Appl. Environ. Microbiol.* 65:1020–28
  69. Gravesse C, Jijakli MH, Lipoivre P. 1998. Study of the exo- $\beta$ -1,3-glucanase activity production by yeast *Pichia anomala* in relation to its antagonistic properties against *Botrytis cinerea* on postharvest apples. *Meded. Fac. Landbouwwet.-Rijksuniv. Gent.* 63:1682–85
  70. Guelldner RC, Reilly CC, Pusey PL, Costello CE, Arrendale RF, et al. 1988. Isolation and identification of iturins as antifungal peptides in biological control of peach brown rot with *Bacillus subtilis*. *J. Agric. Food Chem.* 36:366–70
  71. Guetsky R, Shtienberg D, Elad Y, Dinooor A. 2001. Combining biocontrol agents to reduce the variability of biological control. *Phytopathology* 91:621–27
  72. Guinebretiere MH, Nguyen-The C, Morrison N, Reich M, Nicot P. 2000. Isolation and characterization of antagonists for the biocontrol of the postharvest wound pathogen *Botrytis cinerea* on strawberry fruits. *J. Food Prot.* 63:386–94
  73. Gullino ML, Kuijpers LAM. 1994.



- Social and political implications of managing plant diseases with restricted fungicides in Europe. *Annu. Rev. Phytopathol.* 32:559–79
74. Holmes GJ, Eckert JW. 1999. Sensitivity of *Penicillium digitatum* and *P. italicum* to postharvest citrus fungicides in California. *Phytopathology* 89:716–21
75. Huang Y, Wild BL, Morris C. 1992. Postharvest biological control of *Penicillium digitatum* decay on citrus fruit by *Bacillus pumilus*. *Ann. Appl. Biol.* 120: 367–72
76. Ippolito A, El-Ghaouth A, Wilson CL, Wisniewski M. 2000. Control of postharvest decay of apple fruit by *Aureobasidium pullulans* and induction of defense responses. *Postharvest Biol. Technol.* 19:265–72
77. Ippolito A, Nigro F. 2000. Impact of preharvest application of biological control agents on postharvest diseases of fresh fruits and vegetables. *Crop Prot.* 19:715–23
78. Ippolito A, Nigro F, Romanazzi G, Campanella V. 1998. Field application of *Aureobasidium pullulans* against *Botrytis* storage rot of strawberry. See Ref. 16a, pp. 127–33
79. Janisiewicz W, Yourman L, Roitman J, Mahoney N. 1991. Postharvest control of blue mold and gray mold of apples and pears by dip treatment with pyrrolnitrin, a metabolite of *Pseudomonas cepacia*. *Plant Dis.* 75:490–94
80. Janisiewicz WJ. 1987. Postharvest biological control of blue mold on apples. *Phytopathology* 77:481–85
81. Janisiewicz WJ. 1988. Biocontrol of postharvest diseases of apples with antagonist mixtures. *Phytopathology* 78: 194–98
82. Janisiewicz WJ. 1988. Biological control of diseases of fruits. In *Biocontrol of Plant Diseases*, ed. KG Mukerji, KL Garg, pp. 153–65. Boca Raton, FL: CRC Press
83. Janisiewicz WJ. 1991. Control of postharvest diseases of fruits with biocontrol agents. In *The Biological Control of Plant Diseases*, ed. J Bay-Petersen, pp. 56–68. Taipei, Taiwan: Food Fertil. Technol. Cent. Asian Pac. Reg.
84. Janisiewicz WJ. 1994. Enhancement of biocontrol of blue mold with the nutrient analog 2-Deoxy-D-glucose on apples and pears. *Appl. Environ. Microbiol.* 68:2671–6
85. Janisiewicz WJ. 1996. Ecological diversity, niche overlap, and coexistence of antagonists used in developing mixtures for biocontrol of postharvest diseases of apples. *Phytopathology* 86:473–79
86. Janisiewicz WJ. 1998. Biocontrol of postharvest diseases of temperate fruits. Challenges and opportunities. In *Plant-Microbe Interactions and Biological Control*, ed. GJ Boland, LD Kuykendall, pp. 171–98. New York: Marcel Dekker
87. Janisiewicz WJ, Bors B. 1995. Development of a microbial community of bacterial and yeast antagonists to control wound-invading postharvest pathogens of fruits. *Appl. Environ. Microbiol.* 61: 3261–67
88. Janisiewicz WJ, Conway WS, Glenn DM, Sams CE. 1998. Integrating biological control and calcium treatment for controlling postharvest decay of apples. *HortScience* 33:105–9
89. Janisiewicz WJ, Conway WS, Leverentz B. 1999. Biological control of postharvest decays of apple can prevent growth of *Escherichia coli* O157:H7 in apple wounds. *J. Food Prot.* 62:1372–75
90. Janisiewicz WJ, Jeffers SN. 1997. Efficacy of commercial formulation of two biofungicides for control of blue mold and gray mold of apples in cold storage. *Crop Prot.* 16:629–33
91. Janisiewicz WJ, Korsten L. 2002. Control of postharvest diseases of vegetables by application of microbes. In *Postharvest Physiology of Vegetables*, ed. J Bredt, J Bartz, pp. 543–62. New York: Marcel Dekker. In press

92. Janisiewicz WJ, Marchi. A. 1992. Control of storage rots on various pear cultivars with a saprophytic strain of *Pseudomonas syringae*. *Plant Dis.* 76:555–60
93. Janisiewicz WJ, Peterson DL, Bors R. 1994. Control of storage decay of apples with *Sporobolomyces roseus*. *Plant Dis.* 78:466–70
94. Janisiewicz WJ, Roitman J. 1988. Biological control of blue mold and gray mold on apple and pear with *Pseudomonas cepacia*. *Phytopathology* 78:1697–700
95. Janisiewicz WJ, Tworowski TJ, Kurtzman CP. 2001. Biocontrol potential of *Metchnikowia pulcherrima* strains against blue mold of apple. *Phytopathology* 91:1098–108
96. Janisiewicz WJ, Tworowski TJ, Sharer C. 2000. Characterizing the mechanism of biological control of postharvest diseases on fruits with a simple method to study competition for nutrients. *Phytopathology* 90:1196–200
97. Janisiewicz WJ, Usall J, Bors B. 1992. Nutritional enhancement of biocontrol of blue mold on apples. *Phytopathology* 82:1364–70
98. Jarvis WR. 1962. The infection of strawberry and raspberry fruits by *Botrytis cinerea*. *Ann. Appl. Biol.* 50:569–75
99. Jeffers SN, Wright TS. 1994. Biological control of postharvest diseases of apples: progress and commercial potential. *N. Engl. Fruit Meet.* 100:100–6
100. Jijakli MH, Lepoivre P. 1998. Characterization of an  $\alpha$ - $\beta$ -1,3-glucanase produced by *Pichia anomala* strain K, antagonist of *Botrytis cinerea* on apples. *Phytopathology* 88:335–43
101. Jijakli MH, Lepoivre P, Tossut P, Thonard P. 1993. Biological control of *Botrytis cinerea* and *Penicillium* sp. on post-harvest apples by two antagonistic yeasts. *Meded. Fac. Landbouwwet.-Rijksuniv. Gent.* 58:1349–58
- 101a. Johnson GI, Highley E, Joyce DC, eds. 1998. *Disease Resistance in Fruit*. Canberra: Aust. Cent. Int. Agric. Res.
102. Jones KA, Burges HD. 1998. Technology of formulation and application. See Ref. 22a, pp. 7–30
103. Kanopathipillai VS. 1988. Nonchemical control of *Colletotrichum mussea*, the Anthracnose pathogen of bananas. *Proc. Int. Congr. Plant Pathol.*, 5<sup>th</sup>, p. 422
104. Kanopathipillai VS, Jantan R. 1986. Approach to biological control of anthracnose fruit rot of bananas. In *First Reg. Symp. Biol. Control. Tropics*, ed. MY Hussein, AG Ibrahim, pp. 387–98. Pertanian: Univ. Pertanian
105. Kerr A. 1980. Biological control of crown gall through production of agrocin 84. *Plant Dis.* 64:25–30
106. Koomen I, Jeffries P. 1993. Effects of antagonistic microorganisms on the post-harvest development of *Colletotrichum gloeosporioides* on mango. *Plant Pathol.* 42:230–37
107. Korsten L. 1993. *Biological control of avocado fruit diseases*. PhD thesis. Univ. Pretoria, SA
108. Korsten L, De Jager ES, De Villiers EE, Lourens A, Kotze JM, Wehner FC. 1995. Evaluation of bacterial epiphytes isolated from avocado leaf and fruit surfaces for biological control of avocado postharvest diseases. *Plant Dis.* 79: 1149–56
109. Korsten L, de Villiers EE, de Jager ES, van Harmelen MWS, Heitmann A. 1993. Biological control of litchi fruit diseases. *SA Growers' Assoc. Yearb.* 5:36–40
110. Korsten L, De Villiers EE, Wehner FC, Kotzé JM. 1997. Field sprays of *Bacillus subtilis* and fungicides for control of preharvest fruit diseases of avocado in South Africa. *Plant Dis.* 81:455–59
111. Korsten L, Jeffries P. 2000. Potential for biological control of diseases caused by *Colletotrichum*. In *Colletotrichum Host Specificity, Pathology and Host-Pathogen Interaction*, ed. D Prusky, S Freeman, MB Dickman, pp. 266–95. St. Paul, MN: APS Press
112. Korsten L, De Villiers EE, Wehner FC,

- Kotze JM. 1994. A review of biological control of postharvest diseases of subtropical fruits. In *Postharvest Handling of Tropical Fruits*, ed. BR Champ, E Highley, GL Johnson, pp. 172–85. Aust.: ACIAR
113. Kovach J, Petzoldt R, Harman GE. 2000. Use of honey bees and bumble bees to disseminate *Trichoderma harzianum* 1295-22 to strawberries for *Botrytis* control. *Biol. Control* 18:235–42
  114. Leibinger W, Breuker B, Hahn M, Mengden K. 1997. Control of postharvest pathogens and colonization of the apple surface by antagonistic microorganisms in the field. *Phytopathology* 87:1103–10
  115. Leverentz B, Conway WS, Alavidze Z, Janisiewicz WJ, Fuchs Y, et al. 2001. Examination of bacteriophages as a biocontrol for *Salmonella* on fresh-cut fruit: a model study. *J. Food Prot.* 64:1116–21
  116. Leverentz B, Janisiewicz WJ, Conway WS, Saftner RA. 2001. Effect of combining biocontrol, heat treatment, and MCP-treatment on the reduction of postharvest decay of 'Delicious' apples. *Phytopathology* 91:S55
  117. Leverentz B, Janisiewicz WJ, Conway WS, Saftner RA, Fuchs Y, et al. 2000. Combining yeasts or a bacterial biocontrol agent and heat treatment to reduce postharvest decay of 'Gala' apples. *Postharvest Biol. Technol.* 21:87–94
  118. Lima G, De Curtis F, Castoria R, De Cicco V. 1998. Activity of the yeasts *Cryptococcus laurentii* and *Rhodotorula glutinis* against post-harvest rots on different fruits. *Biocontrol Sci. Technol.* 8: 257–67
  119. Lima G, Ippolito A, Nigro F, Salerno M. 1997. Effectiveness of *Aureobasidium pullulans* and *Candida oleophila* against postharvest strawberry rots. *Postharvest Biol. Technol.* 10:169–78
  120. Limanceau P, Bakker PA, DeKogel WJ, Alabouvette C, Schippers B. 1992. Effect of pseudobactin 358 production by *Pseudomonas putida* WCS358 on suppression of Fusarium wilt of carnations by nonpathogenic *Fusarium oxysporum* Fo47. *Appl. Environ. Microbiol.* 59: 2978–82
  121. Lindow SE, Monier J-M, Leveau JHJ. 2001. Characterizing the microhabitats of bacteria on leaves. *Int. Congr. Mol. Plant-Microbe Interact.*, 10<sup>th</sup>. St. Paul, MN
  122. Mari M, Carati A. 1998. Use of *Saccharomyces cerevisiae* with ethanol in the biological control of grey mould on pome fruits. See Ref. 16a, pp. 85–91
  123. Mari M, Guizzardi M, Pratella GC. 1996. Biological control of gray mold in pears by antagonistic bacteria. *Biol. Control* 7:30–37
  124. Matteson Heidenreich MC, Corral-Garcia MR, Momol EA, Burr TJ. 1997. Russet of apple fruit caused by *Aureobasidium pullulans* and *Rhodotorula glutinis*. *Plant Dis.* 81:339–42
  125. McGuire RC. 1994. Application of *Candida guilliermondii* in commercial citrus coatings for biocontrol of *Penicillium digitatum* on grapefruits. *Biol. Control* 4:1–7
  126. McLaughlin RJ, Wilson CL, Chalutz E, Kurtzman CP, Fett WF, Osman SF. 1990. Characterization and reclassification of yeast used for biological control of postharvest diseases of fruits and vegetables. *Appl. Environ. Microbiol.* 56:3583–86
  127. McLaughlin RJ, Wilson CL, Droby S, Ben-Arie R, Chalutz E. 1992. Biological control of postharvest diseases of grape, peach, and apple with the yeasts *Kloeckera apiculata* and *Candida guilliermondii*. *Plant Dis.* 76:470–73
  128. McLaughlin RJ, Wisniewski ME, Wilson CL, Chalutz E. 1990. Effect of inoculum concentration and salt solutions on biological control of postharvest diseases of apple with *Candida* sp. *Phytopathology* 80:456–61
  129. Mercier J, Arul J, Ponnampalam R, Boulet M. 1993. Induction of 6-methoxymellein and resistance to storage pathogens

- in carrot slices by UV-C. *J. Phytopathol.* 137:44–54
130. Mercier J, Wilson CL. 1994. Colonization of apple wounds by naturally occurring microflora and introduced *Candida oleophila* and their effect on infection by *Botrytis cinerea* during storage. *Biol. Control* 4:138–44
  131. Deleted in proof
  132. Miller WG, Leveau JHJ, Lindow SE. 2000. Improved *gfp* and *inaZ* broad-host-range promoter-probe vectors. *Mol. Plant-Microbe Interact.* 13:1243–50
  - 132a. Moline HE, ed. 1984. *Postharvest Pathology of Fruits and Vegetables: Postharvest Losses in Perishable Crops*. UC, Berkeley: Agric. Exp. Stn.
  133. Nigro F, Finetti Sialer MM, Gallitelli D. 1999. Transformation of *Metschnikowia pulcherrima* 320, biocontrol agent of storage rots, with the green fluorescent protein gene. *J. Plant Pathol.* 81:205–8
  134. Penrose LJ, Nicholls MR, Koffman W. 1987. Failure of UV irradiation to reduce blue mould infections beneficially in an experimental water handling system for pome fruit. *Crop Prot.* 6:399–401
  135. Pierson EA, Weller DM. 1994. Use of mixture of fluorescent pseudomonads to suppress take-all and improve the growth of wheat. *Phytopathology* 84:940–47
  136. Powell KA. 1992. Biocontrol product fermentation, formulation and marketing. See Ref. 175a, pp. 381–87
  137. Prusky D, Bazak M, Ben-Arie R. 1985. Development, persistence, survival and strategies for control of thiabendazole-resistant strains of *Penicillium expansum* on pome fruits. *Phytopathology* 75:877–82
  138. Pusey PL. 1991. Antibiosis as mode of action in postharvest biological control. See Ref. 184a, pp. 127–41
  139. Pusey PL, Wilson CL. 1984. Postharvest biological control of stone fruit brown rot by *Bacillus subtilis*. *Plant Dis.* 68:753–56
  140. Pusey PL, Wilson CL, Hotchkiss MW, Franklin JD. 1986. Compatibility of *Bacillus subtilis* for postharvest control of peach brown rot with commercial fruit waxes, dicloran, and cold-storage conditions. *Plant Dis.* 70:587–90
  141. Qing F, Shiping T. 2000. Postharvest biological control of rhizopus rot of nectarine fruits by *Pichia membranefaciens*. *Plant Dis.* 84:1212–16
  142. Qing F, Shiping T. 2001. Postharvest biological control of grey mold and blue mold on apple by *Cryptococcus albidus* (Saito) Skinner. *Postharvest Biol. Technol.* 21:341–50
  143. Ragsdale NN, Sisler HD. 1994. Social and political implications of managing plant diseases with decreased availability of fungicides in the United States. *Annu. Rev. Phytopathol.* 32:545–57
  144. Raupach GS, Kloepper JW. 1988. Mixtures of plant growth promoting rhizobacteria enhance biological control of multiple cucumber pathogens. *Phytopathology* 88:1158–64
  145. Rishbeth J. 1963. Stump protection against *Fomes annosus*. III. Inoculation with *Peniophora gigantea*. *J. Appl. Biol.* 52:63–77
  146. Rist DL, Rosenberger DA. 1995. A storage decay of apple fruit caused by *Aureobasidium pullulans*. *Plant Dis.* 79:425
  147. Roberts RG. 1990. Postharvest biological control of gray mold of apple by *Cryptococcus laurentii*. *Phytopathology* 80:526–30
  148. Rosenberger DA, Meyer FW. 1981. Postharvest fungicides for apples: development of resistance to benomyl, vinclozolin, and iprodione. *Plant Dis.* 65:1010–13
  149. Schena L, Ippolito A, Zahavi T, Cohen L, Nigro F, Droby S. 1999. Genetic diversity and biocontrol activity of *Aureobasidium pullulans* isolates against postharvest rots. *Postharvest Biol. Technol.* 17:189–99
  150. Schiewe A, Mendgen K. 1992. Identification of antagonists for biological

- control of the post-harvest pathogens *Pezizula malicorticis* and *Nectria galligena* on apples. *J. Phytopathol.* 134:229–37
151. Schisler DA, Slininger PJ, Bothast RJ. 1997. Effects of antagonist cell concentration and two-strain mixtures on biological control of fusarium dry rot of potatoes. *Phytopathology* 87:177–83
  152. Sholberg PL, Marchi A, Bechard J. 1995. Biocontrol of postharvest diseases of apple using *Bacillus* spp. isolated from stored apples. *Can. J. Microbiol.* 41:247–52
  153. Singh V, Deverall BJ. 1984. *Bacillus subtilis* as a control agent against fungal pathogens of citrus fruit. *Trans. Br. Mycol. Soc.* 83:487–90
  154. Slininger PJ, Jackson MA. 1991. Nutritional factors regulating growth and accumulation of phenazine 1-carboxylic acid by *Pseudomonas fluorescens* 2-79. *Appl. Microbiol. Biotechnol.* 37:388–92
  155. Smilanick JL, Denis-Arrue R. 1992. Control of green mold of lemons with *Pseudomonas* species. *Plant Dis.* 76: 481–85
  156. Smilanick JL, Denis-Arrue R, Bosch JR, Gonzalez AR, Henson D, Janisiewicz WJ. 1993. Control of postharvest brown rot of nectarines and peaches by *Pseudomonas* species. *Crop Prot.* 12:513–20
  157. Smilanick JL, Mackey BE, Reese R, Usall J, Margosan DA. 1997. Influence of concentration of soda ash, temperature, and immersion period on the control of postharvest green mold of oranges. *Plant Dis.* 81:379–82
  158. Smilanick JL, Margosan DA, Henson DJ. 1995. Evaluation of heated solutions of sulfur dioxide, ethanol, and hydrogen peroxide to control postharvest green mold of lemons. *Plant Dis.* 79:742–47
  159. Smilanick JL, Margosan DA, Milkota F, Usall J, Michael I. 1999. Control of citrus green mold by carbonate and biocarbonate salts and influence of commercial postharvest practices on their efficacy. *Plant Dis.* 83:139–45
  160. Smolka S. 1992. Methods for in vitro-assessment of pesticide effects on microorganisms of the phyllosphere. *Nachrichtenbl. Dtsch. Pflanzenschutzdienst.* 44: 252–64
  161. Smoot JJ, Melvin CF. 1965. Reduction of citrus decay by hot-water treatment. *Plant Dis. Rep.* 49:463–67
  162. Sobiczewski P, Bryk H. 1996. Biocontrol of *Botrytis cinerea* and *Penicillium expansum* on postharvest apples by antagonistic bacteria. *Int. Conf. Integrated Fruit Prod.*, ed. F Polesny, W Muller, RW Olszak, 19:344–45
  163. Sobiczewski P, Bryk H, Berczynski S. 1996. Evaluation of epiphytic bacteria isolated from apple leaves in the control of postharvest apple diseases. *J. Fruit Ornament. Res.* 4:35–45
  164. Spotts RA. 1984. Environmental modification for control of postharvest decays. See Ref. 132a, pp. 67–72
  165. Spotts RA, Chen PM. 1987. Prestorage heat treatment for control of decay of pear fruit. *Phytopathology* 77:1578–82
  166. Stack JP. 1998. *Postharvest biological control: commercial successes and a model for public and private sector co-operation*. Presented at Int. Congr. Plant Pathol., 7<sup>th</sup>, London. Abstr. 1
  167. Stevens C, Khan VA, Tang AY, Lu J. 1990. The effect of ultraviolet radiation on mold rots and nutrients of stored sweet potatoes. *J. Food Prot.* 53:223–26
  168. Stevens C, Wilson CL, Lu JY, Khan VA, Chalutz E, et al. 1996. Plant hormesis induced by ultraviolet light-C for controlling postharvest diseases of tree fruits. *Crop Prot.* 15:129–34
  169. Sutton JC, Li D, Peng G, Yu H, Zhang P, Valdebenito-Sanhueza RM. 1997. *Gliocladium roseum*, a versatile adversary of *Botrytis cinerea* in crops. *Plant Dis.* 81: 316–28
  170. Takeda F, Janisiewicz WJ. Extending

- strawberry fruit shelf life with pyrrolnitrin. In *From the Strawberry Into 21st Century*, ed. A Dale, JJ Luby, pp. 174–76. Portland, OR: Timber Press
171. Takesako K, Ikai K, Haruna F, Endo M, Shimanaka K, et al. 1991. Aureobasidins, new antifungal antibiotics. Taxonomy, fermentation, isolation, and properties. *J. Antibiot.* 44:919–24
  172. Teixidó N, Usall J, Viñas I. 1999. Efficacy of preharvest and postharvest *Candida sake* biocontrol treatments to prevent blue mould on apples during cold storage. *Int. J. Food Microbiol.* 50:203–10
  173. Teixidó N, Viñas I, Usall J, Magan N. 1998. Control of blue mold of apples by preharvest application of *Candida sake* grown in media with different water activity. *Phytopathology* 88:960–64
  174. Teixidó N, Viñas I, Usall J, Sanchis V, Magan N. 1998. Ecophysiological responses of the biocontrol yeast *Candida sake* to water, temperature and pH stress. *J. Appl. Microbiol.* 84:192–200
  175. Testoni A, Aloï C, Mocioni M, Gullino GL. 1993. Biological control of *Botrytis* rot of kiwi fruit. *Bull. Biol. Control Foliar Post-Harvest Dis.* 16:95–98
  - 175a. Tjamos C, Papavizas GC, Cook RJ, eds. 1992. *Biological Control of Plant Diseases: Progress and Challenges for the Future*. Adv. Res. Works. New York: Plenum
  176. Tronsmo A, Dennis C. 1977. The use of *Trichoderma* species to control strawberry fruit rots. *Neth. J. Plant Pathol.* 83:449–55
  177. Tukey B. 1993. Overview of ozone use at Snokist Growers. *Tree Fruit Postharvest J.* 4:14–15
  178. Usall J, Teixidó N, Fons E, Viñas I. 2000. Biological control of blue mould on apple by a strain of *Candida sake* under several controlled atmosphere conditions. *Int. J. Food Microbiol.* 58:83–92
  179. Utkhede RS, Sholberg PL. 1986. In vitro inhibition of plant pathogens by *Bacillus subtilis* and *Enterobacter aerogenes* and in vivo control of two postharvest cherry diseases. *Can. J. Microbiol.* 32:963–67
  180. Viñas I, Usall J, Sanchis V. 1991. Tolerance of *Penicillium expansum* to postharvest fungicide treatments in apple packinghouses in Lerida (Spain). *Mycopathologia* 113:15–18
  181. Vishniac HS, Anderson JA, Filonow AB. 1997. Assimilation of volatiles from ripe apples by *Sporidiobolus salmonicolor* and *Tilletiopsis washingtonensis*. *Antonie van Leeuwenhoek J. Microbiol. Serol.* 72:201–7
  182. Weller D. 1988. Biological control of soilborne plant pathogens in the rhizosphere with bacteria. *Annu. Rev. Phytopathol.* 26:379–407
  183. Wilson C, Pusey PL. 1985. Potential for biological control of postharvest plant diseases. *Plant Dis.* 69:375–78
  184. Wilson CL, Chalutz E. 1989. Postharvest biological control of *Penicillium* rots of citrus with antagonistic yeasts and bacteria. *Sci. Hortic.* 40:105–12
  - 184a. Wilson CL, Chalutz E, eds. 1991. *Biological Control of Postharvest Diseases of Fruits and Vegetables, Workshop Proc.* Washington, DC: US GPO
  185. Wilson CL, El Ghaouth A, Chalutz E, Droby S, Stevens C, et al. 1994. Potential of induced resistance to control postharvest diseases of fruits and vegetables. *Plant Dis.* 78:837–43
  186. Wilson CL, Wisniewski ME. 1989. Biological control of postharvest diseases of fruits and vegetables: an emerging technology. *Annu. Rev. Phytopathol.* 27: 425–41
  187. Wilson CL, Wisniewski ME, Droby E, Chalutz E. 1993. A selection strategy for microbial antagonists to control postharvest diseases of fruits and vegetables. *Sci. Hortic.* 53:183–89
  188. Wilson CL, El Ghaouth A. 1993. Multifaceted biological control of postharvest diseases of fruits and vegetables. *Proc. Beltsville Symp.* 18<sup>th</sup>, ARS USDA,

- May 2–6, ed. RD Lumsden, JL Vaughn. Washington, DC: Am. Chem. Soc.
189. Wilson M, Lindow S. 1994. Ecological similarity and coexistence of epiphytic ice-nucleating ( $\text{Ice}^+$ ) *Pseudomonas syringae* strains and a non-ice-nucleating ( $\text{Ice}^-$ ) biological control agent. *Appl. Environ. Microbiol.* 60:3128–37
190. Wisniewski M, Biles C, Droby S, McLaughlin R, Wilson C, Chalutz E. 1991. Mode of action of the postharvest biocontrol yeast, *Pichia guilliermondii*. 1. Characterization of attachment to *Botrytis cinerea*. *Physiol. Mol. Plant Pathol.* 39:245–58
191. Wisniewski M, Droby S, Chalutz E, Eilam Y. 1995. Effects of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  on *Botrytis cinerea* and *Penicillium expansum* *in vitro* and on the biocontrol activity of *Candida oleophila*. *Plant Pathol.* 44:1016–24
192. Wisniewski M, Wilson C, Chalutz E, Hershberger W. 1988. *Biological control of postharvest diseases of fruit: inhibition of Botrytis rot on apple by an antagonistic yeast*. Presented at Annu. Meet. Electron Microsc. Soc. Am., 46<sup>th</sup>, San Francisco
193. Wisniewski M, Wilson C, Hershberger W. 1989. Characterization of inhibition of *Rhizopus stolonifer* germination and growth by *Enterobacter cloacae*. *Can. J. Bot.* 67:2317–23
194. Yehuda H, Droby S, Wisniewski M, Goldway M. 2001. A transformation system for the biocontrol yeast, *Candida oleophila*, based on hygromycin B resistance. *Curr. Genet.* 40:276–81
195. Zahavi T, Cohen L, Weiss B, Schena L, Daus A, et al. 2000. Biological control of *Botrytis*, *Aspergillus* and *Rhizopus* rots on table and wine grapes in Israel. *Postharvest Biol. Technol.* 20:115–24
196. Zhou T, Northover J, Schneider KE. 1999. Biological control of postharvest diseases of peach with phyllosphere isolates of *Pseudomonas syringae*. *Can. J. Plant Pathol.* 21:375–81



## CONTENTS

---

FRONTISPIECE	xii
ONE FOOT IN THE FURROW: IMPLICATIONS TO ONE'S CAREER IN PLANT PATHOLOGY, <i>D. E. Mathre</i>	1
EVOLUTIONARY ECOLOGY OF PLANT DISEASES IN NATURAL ECOSYSTEMS, <i>Gregory S. Gilbert</i>	13
MAKING AN ALLY FROM AN ENEMY: PLANT VIROLOGY AND THE NEW AGRICULTURE, <i>Gregory P. Pogue, John A. Lindbo, Stephen J. Garger, and Wayne P. Fitzmaurice</i>	45
STRIPE RUST OF WHEAT AND BARLEY IN NORTH AMERICA: A RETROSPECTIVE HISTORICAL REVIEW, <i>Roland F. Line</i>	75
VIRAL SEQUENCES INTEGRATED INTO PLANT GENOMES, <i>Glyn Harper, Roger Hull, Ben Lockhart, and Neil Olszewski</i>	119
MOLECULAR BASIS OF RECOGNITION BETWEEN <i>PHYTOPHTHORA</i> PATHOGENS AND THEIR HOSTS, <i>Brett M. Tyler</i>	137
COMPARATIVE GENOMIC ANALYSIS OF PLANT-ASSOCIATED BACTERIA, <i>M. A. Van Sluys, C. B. Monteiro-Vitorello, L. E. A. Camargo, C. F. M. Menck, A. C. R. da Silva, J. A. Ferro, M. C. Oliveira, J. C. Setubal, J. P. Kitajima, and A. J. Simpson</i>	169
GENE EXPRESSION IN NEMATODE FEEDING SITES, <i>Godelieve Gheysen and Carmen Fenoll</i>	191
PHYTOCHEMICAL BASED STRATEGIES FOR NEMATODE CONTROL, <i>David J. Chitwood</i>	221
HOST-SELECTIVE TOXINS AND AVIRULENCE DETERMINANTS: WHAT'S IN A NAME? <i>Thomas J. Wolpert, Larry D. Dunkle, and Lynda M. Ciuffetti</i>	251
TOBACCO MOSAIC VIRUS ASSEMBLY AND DISASSEMBLY: DETERMINANTS IN PATHOGENICITY AND RESISTANCE, <i>James N. Culver</i>	287
MICROBIAL POPULATIONS RESPONSIBLE FOR SPECIFIC SOIL SUPPRESSIVENESS TO PLANT PATHOGENS, <i>David M. Weller, Jos M. Raaijmakers, Brian B. McSpadden Gardener, and Linda S. Thomashow</i>	309



PATHOGEN POPULATION GENETICS, EVOLUTIONARY POTENTIAL, AND DURABLE RESISTANCE, <i>Bruce A. McDonald and Celeste Linde</i>	349
USE OF MULTILINE CULTIVARS AND CULTIVAR MIXTURES FOR DISEASE MANAGEMENT, <i>C. C. Mundt</i>	381
BIOLOGICAL CONTROL OF POSTHARVEST DISEASES OF FRUITS, <i>Wojciech J. Janisiewicz and Lise Korsten</i>	411
ANTIBIOTIC USE IN PLANT AGRICULTURE, <i>Patricia S. McManus,</i> <i>Virginia O. Stockwell, George W. Sundin, and Alan L. Jones</i>	443
RISK ASSESSMENT OF VIRUS-RESISTANT TRANSGENIC PLANTS, <i>Mark Tepfer</i>	467
INDEXES	
Subject Index	493
Cumulative Index of Contributing Authors, Volumes 31–40	521
Cumulative Index of Chapter Titles, Volumes 31–40	525
ERRATA	
An online log of corrections to <i>Annual Review of Phytopathology</i> chapters (if any, 1997 to the present) may be found at <a href="http://phyto.annualreviews.org/">http://phyto.annualreviews.org/</a>	